

*profunda* (Say), and *Triodopsis multilineata* (Say) occasionally exhibit a transitory, dome-like swelling between the dorsal tentacles. This phenomenon, which appears most frequently during periods of increased mating activity in the teraria, has been observed only in adults—usually in animals with swollen and prominent genital areas. Although sometimes observed immediately prior to and even (in *T. multilineata*) during courtship maneuvers, the swelling is most prominently displayed by animals that sit in solitude and remain almost motionless for considerable periods. All observations point to the conclusion that this display usually precedes sexual encounter and courtship.

Of possible explanations, the simplest is that the swelling is a mechanical result of internal pressures coincident with sexual arousal and that it has no other particular significance. A more attractive possibility is that the swelling is a specialized emitter or receptor site involved in chemical attraction or detection of potential mates.

(Several slides of living snails exhibiting the abnormality were shown.) Morrison: "I don't think you can consider this a chemical receptor. It seems more likely to be some generic abnormality."

#### FRESHWATER MOLLUSKS OF THE HUDSON BAY WATERSHED, DISTRIBUTION PATTERNS AND DETERMINANT INFLUENCES. Arthur H. Clarke, Jr., National Museums of Canada, Ottawa, Ontario, Canada.

##### (Abstract)

Just north of the Mississippi-Missouri and St. Lawrence basins lies a vast area of partially interconnected river systems, the Hudson Bay Watershed. About 100 species and subspecies of freshwater mollusks occur here, and all major North American families except Pleuroceridae are represented.

Based on distribution, three general groups can be distinguished: (I) endemic elements (about 10% of the total); (II) holarctic elements (about 15%); and (III) southern elements (about 75%). Group I contains several taxa which appear to have evolved since the Pleistocene (e.g., *Valvata sinerea ontariensis* and *Stagnicola aretina*). Group II includes species which penetrate farther into the Canadian arctic than any others, i.e., to Baffin Island and Victoria Island. Members of this group include *Lymnaea stagnalis*, *Stagnicola palustris*, several sphaeriids, etc. Group III is composed principally of species which reach their northern limits within this region, and it is sometimes possible to correlate these limits with biological and geological factors and with isotherms.

In the Unionidae, for example, several species apparently invaded the Hudson Bay Watershed during postglacial confluence from the Mississippi-Missouri Basin by way of the Red River. Some of these (*Fusconaia flava*, *Quadrula quadrula*, *Crenodonta plicata*, *Lasmigona costata*, and *Proptera nitata*) have not penetrated beyond southern Manitoba; others (*Ligumia recta*, *Lampsilis ovata ventricosa*, and *Siphonaria undulatus*) have spread into northern and western Manitoba or to eastern Saskatchewan; and one (*Lasmigona complanata*) has spread even through central Alberta to the Lake Athabasca drainage. All range expansions in this group have been northward and west-

ward, never eastward into the soft-water habitats of the Precambrian Shield. The varying degrees of success in penetrating colder regions may be related to threshold spawning requirements. In only two cases (*Ligumia recta* and *Siphonaria undulatus*) does absence of the known fish host appear to be limiting.

Other Unionidae have used alternate invasion routes. *Elliptio complanata* has penetrated the Hudson Bay Watershed only from the St. Lawrence System and is confined to that part of Ontario and Quebec east of Lake Nipigon and south of James Bay. It is uncommon or rare in this region, however, and competition with the abundant and nearby ubiquitous *Lampsilis radiata* siliqua may be a limiting factor. Two other species, *Lasmigona compressa* and *Anodontoides ferussacianus*, appear to have used dual invasion routes, one from Lake Superior through Lake Nipigon and Long Lac and one from the Mississippi-Missouri Basin through the Red River, but the distribution of neither species in the Hudson Bay Watershed appears to be correlated with bedrock geology or with temperature.

*Anodonta grandis* and *L. radiata* siliqua may have utilized multiple invasion routes. Both species occur throughout the entire boreal forest region even including the Mackenzie River north of Great Slave Lake. Summer water temperature data are desirable but unavailable. The northern limits of both species fall within the zone bounded by the 55° and 60°F. July average daily temperature isotherms, however, and are probably indirectly related to these average values.

(A projected map of Canada, of the geology of the region and slides of species followed by distribution maps illustrated Dr. Clarke's paper.)  
President Burch: "This paper represents a great amount of painstaking work!" \* \* \*

With this, opening day's papers were brought to an end and the three hours before the scheduled evening assembly afforded the first opportunity to locate and sample the cuisine of one or another of the fine restaurants so plentiful in New Orleans. In this, home of the po' boy sandwich and where grits and red-eye gravy are as staple as bread and butter, the Northerners were intrigued by the variety of unfamiliar dishes. Perhaps most enjoyed was the abundance of fish and crustacea, never better than here, short hours away from their watery home.

At eight o'clock began an innovation which had been billed as Shell Club Night. Each of the thirty-two local shell clubs affiliated with the AMU had been invited to participate by sending delegates and an exhibit, a report of club activities, or other offerings of an instructive or entertaining nature. Mr. and Mrs. Harvey Meyer acted as joint chairmen of the evening, doing a masterful job of correlating the varied material into a smooth-running and interesting program.

#### SHELL CLUB NIGHT

Spokesmen for the North Carolina Shell Club were Carl Withrow and James Wadsworth; it covers, said they, probably the largest geographical area of any club. Three meetings are held on the coast each year, the fourth inland.